



POSTER PRESENTATION

Open Access

Immune response characterization in HIV/HCV co-infected patients of medicine tropical foundation

Adriana Malheiro^{1,3*}, Liziara Silva Fraport^{1,3}, Flamir Victoria², Kátia Luz Torres¹, João Paulo Diniz Pimentel¹, Andrea Tarragô¹, Laura Patricia Viana Maia^{1,3}, Felicien Vásquez¹, José Eduardo Levi⁴, Marilu Victoria²

From 16th International Symposium on HIV and Emerging Infectious Diseases
Marseille, France. 24-26 March 2010

Background

The epidemiology of co-infection of human immunodeficiency virus and hepatitis C virus (HIV/HCV) is around 30 to 60%. Approximately one third of HIV infected shows C hepatitis, with a high rate in hemophiliacs and drug users. Recent publications demonstrated that HIV positive patients co-infection with HCV have a co-factor to develop AIDS. The purpose of this study was evaluate the cellular and humoral immune response and cytokines in HIV/HCV co-infected patients in Foundation of Tropical Medicine of Amazonas.

Methods

After consent term assignature, the population of T lymphocytes CD4⁺ and CD8⁺ was analyzed in the whole blood by flow cytometry and a blood sample was take to measure the serum concentration of inflammatory cytokines (interleucine – IL - 6, 8 and tumoral necrosis factor alpha-TNF- α), cytokines of T_H1 (IL-12, Interferon gamma-IFN- γ) cytokines T_H2 (IL-4) and suppression cytokine (IL-10) using ELISA BD OptEIA[®] kit.

Results

As for CD4⁺T cells 72.2% had < 500 cls/mm³ with a median of 271 cls/mm³, on the T CD8⁺ 88.9% had \geq 215 cls/mm³ with a median of 794.5 cls/mm³. The ratio CD4⁺/CD8⁺ was 0.32 cls/mm³. When the dose cytokines IL-4, IL-6, IL-8, IL-10, IL-12 and IFN- γ in the patients found that only the IL-6 ($p = < 0.001$) showed statistical significance especially when correlated to the logarithm of the HCV viral load (0.031).

Discussion

The results found in this study, despite the low prevalence, have annual growth of co-infection due to improvement in the research of hepatitis C in patients with HIV and the IL-6 cytokine was important marked of inflammation in this studied population.

Financial support

FHEMOAM, FAPEAM.

Author details

¹Fundação de Hematologia e Hemoterapia do Amazonas, Manaus, Brazil.

²Fundação de Medicina Tropical do Amazonas, Manaus, Brazil. ³Universidade Federal do Amazonas, Manaus, Brazil. ⁴Universidade Estadual de São Paulo, São Paulo, Brazil.

Published: 11 May 2010

doi:10.1186/1742-4690-7-S1-P80

Cite this article as: Malheiro et al.: Immune response characterization in HIV/HCV co-infected patients of medicine tropical foundation. *Retrovirology* 2010 **7**(Suppl 1):P80.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: elisadleon@yahoo.com.br

¹Fundação de Hematologia e Hemoterapia do Amazonas, Manaus, Brazil